



King's Research Portal

DOI:

[10.1007/s00204-015-1593-7](https://doi.org/10.1007/s00204-015-1593-7)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Shi, Q., Haenen, G. R., Maas, L., Arlt, V. M., Spina, D., Vasquez, Y. R., Moonen, E., Veith, C., Van Schooten, F. J., & Godschalk, R. W. L. (2015). Inflammation-associated extracellular -glucuronidase alters cellular responses to the chemical carcinogen benzo[a]pyrene. *Archives of Toxicology*, 1-13. <https://doi.org/10.1007/s00204-015-1593-7>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Inflammation associated extracellular β -glucuronidase alters cellular responses to the chemical carcinogen benzo[a]pyrene

Shi Q¹, Haenen GR¹, Maas L¹, Arlt VM^{2,3}, Spina D⁴, Rizzo Vasquez Y⁴, Moonen E¹, Veith C¹, Van Schooten FJ¹, Godschalk RWL¹

¹Department of Pharmacology & Toxicology, NUTRIM School of Nutrition and Translational Research in Metabolism, Maastricht University, PO Box 616, 6200 MD, Maastricht, The Netherlands

²Analytical and Environmental Sciences Division, King's College London, MRC-PHE Centre for Environmental & Health, 150 Stamford Street, London SE1 9NH, United Kingdom

³NIHR Health Protection Research Unit in Health Impact of Environmental Hazards at King's College London in partnership with Public Health England, 150 Stamford Street, London SE1 9NH, United Kingdom

⁴Sackler Institute of Pulmonary Pharmacology, Institute of Pharmaceutical Science, King's College London, 150 Stamford Street, London SE1 9NH, United Kingdom

To whom correspondence should be addressed. Tel: +31 43 3881104, Fax: +31 43 3884146, Email:

R.Godschalk@maastrichtuniversity.nl

Abstract

Neutrophils infiltrate tissues during inflammation and when activated, they release β -glucuronidase. Since inflammation is associated with carcinogenesis, we investigated how extracellular β -glucuronidase changed the *in vitro* cellular response to the chemical carcinogen benzo(a)pyrene (B[a]P). For this we exposed human liver (HepG2) and lung (A549) cells to B[a]P in the presence or absence of β -glucuronidase. β -Glucuronidase reduced B[a]P-induced expression of *CYP1A1* and *CYP1B1* at 6 hours after exposure, which did not depend on β -glucuronidase activity, because the inhibitor D-saccharic acid 1,4-lactone monohydrate did not antagonize the effect of β -glucuronidase. On the other hand, the inhibitory effect of β -glucuronidase on CYP expression was dependent on signaling via the insulin like growth factor receptor (IGF2R, a known receptor for β -glucuronidase), because co-incubation with the IGF2R inhibitor mannose-6-phosphate (M6P) completely abolished the effect of β -glucuronidase. Extracellular β -glucuronidase also reduced the formation of several B[a]P metabolites and B[a]P-DNA adducts. Interestingly, at 24 hours of exposure, β -glucuronidase significantly enhanced *CYP* expression, probably because β -glucuronidase de-glucuronidated B[a]P metabolites, which continued to trigger the aryl hydrocarbon receptor (Ah-receptor) and induced expression of *CYP1A1* (in both cell lines) and *CYP1B1* (in A549 only). Consequently, significantly higher concentrations of B[a]P metabolites and DNA adducts were found in β -glucuronidase-treated cells at 24 hrs. DNA adduct levels peaked at 48 hours in cells that were exposed to B[a]P and treated with β -glucuronidase. Overall, these data show that β -glucuronidase alters the cellular response to B[a]P and ultimately enhances B[a]P-induced DNA adduct levels.

Key words: Benzo[a]pyrene, Inflammation, β -glucuronidase, Cytochrome P450 1A1, carcinogen metabolism, IGF2R, DNA adducts

Introduction

Chronic inflammation is causally associated with cancer development and therefore, inflammation was considered as the seventh hallmark feature of cancer (Colotta et al. 2009; Shacter and Weitzman 2002). This is illustrated for instance by the relatively high incidence of lung cancer in chronic obstructive pulmonary disease (COPD) patients (Young et al. 2009). Recent studies have pointed out that polymorphonuclear neutrophils (PMN), which are recruited at the site of inflammation play an important role in the initiation and progression of cancer (Fridlender and Albelda 2012; Knaapen et al. 2006). Inhalation of complex air pollutants like tobacco smoke and/or fine particles may result in pulmonary inflammation, which generates reactive oxygen/nitrogen species (ROS/RNS) that can damage lung tissue (van Berlo et al. 2010). At the same time it has been shown that under inflammatory conditions, PMN enhance the mutagenic potential of chemical carcinogens (Borm et al. 1997; Van Schooten et al. 2004). This is partly explained by the action of the PMN-derived enzyme myeloperoxidase (MPO), that can metabolically activate carcinogens and inhibit DNA repair, leading to higher levels of carcinogen-DNA adducts (Gungor et al. 2010a; Gungor et al. 2010b; Gungor et al. 2010c). However, there is still little data available on other factors that are released by the relatively high number of PMN during chronic inflammation, including β -glucuronidase, and how these influence the cellular response to chemical carcinogens (Basinska and Florianczyk 2003).

Inhalatory exposure to air pollutants like particles often results in a combination of an inflammatory response in the presence of genotoxic agents, like polycyclic aromatic hydrocarbons (PAHs) (Hoffmann and Hoffmann 1997). PAHs, including benzo[a]pyrene (B[a]P) have gained much attention, because they are abundantly present in the environment (Uppstad et al. 2010) and they have serious adverse genotoxic effects. B[a]P becomes mutagenic and carcinogenic after bioactivation by enzymes, including cytochrome P450 (CYP) and epoxide hydrolase (EH) (Stiborova et al. 2014). It is converted into various metabolites, including oxides, phenols, diols, diol-epoxides, quinones, and radical cations (Shimada and Guengerich 2006). The best studied metabolite is B[a]P-diol epoxide (BPDE) (**Figure 1**). The first steps is the conversion of B[a]P by the microsomal NADPH-dependent cytochrome P450 isoforms 1A1 (CYP1A1) and 1B1 (CYP1B1) to yield the B[a]P-7,8-oxide, B[a]P-9,10-oxide or 3-hydroxy-B[a]P (3-OH-B[a]P) (Krais et al. 2015; Wohak et al. 2014). Subsequently, both B[a]P-7,8-oxide and B[a]P-9,10-oxide can be hydrated by microsomal EH to yield the corresponding B[a]P-7,8-*trans*-dihydrodiols (B[a]P-7,8-diol) and B[a]P-9,10-*trans*-dihydrodiols (B[a]P-9,10-diol). B[a]P-7,8-diol is further metabolized to the ultimate carcinogen benzo[a]pyrene-7,8-diol-9,10-epoxide (BPDE), which is known as a reactive derivative of B[a]P that can

covalently bind to DNA to form adducts preferentially at guanine residues (*e.g.* 10-(deoxyguanosin- N^2 -yl)-7,8,9-trihydroxy-7,8,9,10-tetrahydro benzo[a]pyrene; dG- N^2 -BPDE) (Arlt et al. 2008; Kim et al. 1998; Wohak et al. 2014). It is generally accepted that the formation of such DNA adducts leads to mutations relevant for carcinogenesis (Kucab et al. 2015). In addition, the reactive metabolites can be conjugated by an important phase II detoxification enzyme UDP-glucuronosyltransferases (UGTs) that leads to glucuronides which are not mutagenic or carcinogenic (Shimoi and Nakayama 2005; Shimoi et al. 2001). Indeed, UGT1A6 was found to glucuronate a range of B[a]P metabolites including 3-OH-B[a]P, B[a]P-9,10-diol and the pro-carcinogen B[a]P-7,8-diol (Jin et al. 1993; Trushin et al. 2012; Zheng et al. 2002).

In order to elucidate the role of β -glucuronidase in inflammatory disease and its effect on chemically-induced cancer, it is crucial to understand how β -glucuronidase changes the cellular response toward B[a]P. In a recent study, it was found that lipopolysaccharide (LPS) treatment increased B[a]P-induced DNA adduct levels in lung and liver tissues of B[a]P inhalatory exposed mice (Arlt et al. 2015). To explain this result we hypothesise that the LPS induced an inflammatory response resulting in the release of β -glucuronidase that hydrolysed glucuronidated B[a]P metabolites and thus reverse the protective effect of glucuronidation, and subsequently enhanced the binding of B[a]P metabolites to DNA. To test our hypothesis, we studied the impact of extracellular β -glucuronidase on the formation of B[a]P metabolites (3-OH-B[a]P, B[a]P-9,10-diol and B[a]P-7,8-diol), gene expression of some enzymes pivotal in B[a]P metabolism (including *CYP1A1*, *CYP1B1* and *UGT1A6*) and DNA adduct formation in human liver cell line (HepG2) and human lung cell line (A549).

Materials and Methods

Mouse lung and liver tissues

All procedures performed in the study involving animal experiments were conducted at King's College London under license in accordance with the institutional ethics committee on the protocols approved by the Home Office under 'The Animals (Scientific Procedures) Act (1986)'. Mice were divided into four groups (n=3 in each group): A) control group; mice were nasally instilled with saline. After 24 hours, mice were intratracheally instilled with tricaprylin. B) the lipopolysaccharide (LPS) treatment group; each mouse was nasally instilled with 20 μ g LPS (dissolved in saline). After 24 hours, mice were intratracheally instilled with tricaprylin. C) the B[a]P-treated group;

mice were nasally instilled with saline and intratracheally instilled with 0.5 mg B[a]P (dissolved in 25 μ l tricaprylin) after 24 hours. D) the B[a]P- and LPS-treated group; each mouse was nasally instilled with 20 μ g of LPS. After 24 hours, mice were intratracheally instilled with 0.5 mg B[a]P. All mice were sacrificed at 48 hours after the intratracheal exposure. The collection of bronchoalveolar lavage fluid (BAL fluid) and the isolation of cytosolic fractions from lung and liver was performed as described in Arlt *et al.* (2015).

Cell lines and cell treatment

Human liver hepatocellular carcinoma HepG2 cells and human epithelial lung adenoma carcinoma A549 cells were obtained from the American Tissue Culture Collection. HepG2 cells were cultured in Minimum Essential Medium (MEM) plus glutamax containing 10% (v/v) foetal calf serum (FCS, Gibco invitrogen, Breda, The Netherlands), 1% (v/v) Sodium Pyruvate, 1% (v/v) penicillin/streptomycin (Sigma, Zwijndrecht, The Netherlands) and 1% (v/v) non-essential amino acids (Sigma). A549 cells were cultured in RPMI (Sigma) supplemented with 5% (v/v) FCS and 1% (v/v) penicillin/streptomycin. All cells were cultured under humidified atmosphere containing 5% CO₂ at 37°C. Cell passages between 20 and 29 were used for experiments. All chemicals were purchased from Sigma-Aldrich unless stated otherwise.

Cells with 80% confluency were exposed to 1 μ M B[a]P in the presence or absence of β -glucuronidase (4 U/ml) for 6, 24 or 48 hours. B[a]P was dissolved in dimethylsulfoxide (DMSO) and added to the medium with a final DMSO concentration of 0.5% (v/v). β -glucuronidase was dissolved in 0.1 M sodium-acetate buffer (pH 5.5). Mannose-6-phosphate (M6P) was dissolved in 0.1 M sodium-acetate buffer (pH 5.5) and the final concentration in cell culture is 100 μ M. D-saccharic acid 1,4-lactone monohydrate was dissolved in 0.1 M sodium-acetate buffer (pH 5.5) the final concentration in cell culture is 100 μ M. Before incubation, all cell media were adjusted to pH 5.5 by using 1 M HCl in order to mimic the micro-environment of inflammation. After incubation, the medium and cells were stored at -20°C until further analysis. Experiments were performed at least with three replicates in three independent cultures.

Measurement of β -glucuronidase activity

β -Glucuronidase (Helix pomatia type H5, $\geq 400,000$ Units/g solid) activity was determined by fluorometrically monitoring the hydrolysis of 4-methylumbelliferyl- β -d-glucuronide (4MUGIA) according to the method described by Bartholome *et al* (Bartholome et al. 2010) with some modifications. Briefly, the reaction mixture in a total volume of 140 μ l contained 0.1 M sodium-acetate buffer (pH 5.5), 2 mM 4MUGIA and sample containing β -glucuronidase. The reaction was initiated by mixing 4MUGIA and the sample. The hydrolysis of 4MUGIA was measured in a thermostated plate reader (Spectra max m2, MDS, CA) at 37°C and 320/460 nm excitation/ emission wavelengths. A standard curve of β -glucuronidase (0.4-400 Units/ml (U/ml)) was generated to quantitate formation of fluorescence in the presence of 4MUGIA. D-saccharic acid 1,4-lactone monohydrate, a β -glucuronidase inhibitor, was used to inhibit β -glucuronidase activity.

For testing the interaction between B[a]P and β -glucuronidase, we used similar condition as mentioned above. A total volume of 140 μ l contained 0.1 M sodium-acetate buffer (pH 5.5), different concentration of 4MUGIA (e.g. 500, 250, 100, 50, 10 and 1 μ M) and 4 U/ml β -glucuronidase with additional 1 μ l of 200 μ M B[a]P or 1 μ l DMSO. The measurement of fluorescence was performed for 10 hours at 37°C.

HPLC fluorescence analysis of B[a]P and B[a]P metabolites

B[a]P and its metabolites were extracted from 5 ml cell medium by mixing with 1 ml ethylacetate for 20 mins and followed by centrifugation (10 min, 980g). The top layer was transferred to a new tube. This procedure was repeated twice. The top layers were evaporated under nitrogen and the residue was redissolved in 0.5 ml methanol (Biosolve Chemicals, Valkenswaard, The Netherlands). Samples were subsequently analyzed by HPLC-FD using a Gynkotek P580A HPLC system (Separations Analytical Instruments, Hendrik-Ido-Ambacht, The Netherlands) consisting of a Spark SP830 autosampler (Spark Holland, Emmen, The Netherlands) and a Perkin Elmer LS-30 programmable fluorescence detector (Perkin Elmer, Foster City, CA, USA) operated at excitation/emission wavelengths 257/ >350 nm. The samples were injected onto a Hypersil 5 μ m ODS HPLC column (250 mm \times 3 mm) (Supelco 54933, Bellefonte, PA, USA) with a flow rate of 0.5 ml/min. Separation was performed using a mixture of two mobile phases: A (100 % methanol) and B (40 % methanol in water) in the following multi-step gradient conditions: 0-5 min,

30/70 (A/B, v/v); 5-30 min, gradient from 30/70 (A/B, v/v) to 90/10 (A/B, v/v); 30-35 min, 90/10 (A/B, v/v); 35-37 min, gradient from 90/10 (A/B, v/v) to 30/70 (A/B, v/v); 37-40 min, 30/70 (A/B, v/v). For quantitation of the specific metabolites, a standard mix which contained 50 ng/ml B[a]P-9,10-diol 50 ng/ml B[a]P-7,8-diol and 50 ng/ml 3-OH-B[a]P (Midwest Research Institute, Kansas City, MO, USA) were injected and the area of each metabolite peak in the chromatogram was determined.

Quantitative real-time PCR

Gene expression levels were measured by quantitative real-time reverse transcriptase-PCR (RT-qPCR) using a MyiQ Single Colour real-time PCR detection system (BioRad, Veenendaal, The Netherlands). Total RNA was isolated and purified by using the RNeasy® Mini Kit (Qiagen Westburg, Leusden, the Netherlands) in combination with DNase treatment (Qiagen). cDNA was generated from 500 ng total RNA by using the iScript™ cDNA synthesis kit protocol (BioRad). Primers were purchased from Operon (Leiden, The Netherlands) for the following genes: *β-actin*, *CYP1A1*, *CYP1B1* and *UGT1A6* (see (Schults et al. 2014)). The reaction contained SYBR® Green Supermix (BIO-RAD), 5 µl (40 times diluted) cDNA and 0.3 µM primers in a total volume of 25 µl. PCR was conducted under the following condition: denaturation at 95°C for 3 min, followed by 40 cycles of 95°C for 10 sec and 55°C for 45 sec. All PCR reactions included a cDNA dilution curve to assess PCR efficiency and all reactions were followed by a melt curve (55-95°C). Data were analyzed by using MyiQ Software system (BioRad) and the amount of target cDNA in each sample was determined by a fractional PCR threshold cycle number (Ct-value) and compared to the corresponding Ct-value for the housekeeping gene *β-actin*. The relative gene expression level for each gene was calculated by using the $2^{-\Delta\Delta C_t}$ method (Livak and Schmittgen 2001).

³²P-postlabelling of B[a]P-DNA adducts

DNA harvested from cells was isolated using a phenol-chloroform-isoamylalcohol extraction procedure as described by Schults *et al.* (Schults et al. 2013). Briefly, after incubation, cells were resuspended in 450 µl lysis buffer [10 mM Tris, 10 mM TEMPO, 1 mM EDTA and 1% (w/v) sodium dodecyl sulfate (SDS); pH 8] and incubated with

proteinase K (10 µg/ml) at 37°C overnight. The mixture was extracted with 1 volume Tris-saturated phenol, 1 volume Tris-saturated phenol-chloroform-isoamyl alcohol (25:24:1 by volume) and 1 volume chloroform-isoamyl alcohol (24:1, v/v). The DNA was precipitated with 1/30 volume 3 M NaAc pH 5.2 and 2 volumes of cold 100% ethanol. Precipitated DNA was washed with 70% ethanol and dried under nitrogen. The DNA was dissolved in 2 mM Tris (pH 8.0) with final concentration 0.5 µg/µl.

DNA digestion and ³²P-postlabelling were performed as described by Van Schooten *et al* (Van Schooten et al. 1997). In short, DNA samples (10 µg) were digested with micrococcal nuclease (Sigma) (0.25 U/µl) and spleen phosphodiesterase (Sigma) (2 µg/µl) for 4 hours at 37°C in a total volume of 9.5 µl. For DNA adduct enrichment, samples were treated with nuclease P1 (Sigma) (2.5 µg/µl) at 37°C for 30 min. The nuclease P1 reaction was terminated by addition of 1 µl 1 M Tris (pH 9.6). DNA adducts were subsequently labelled with [γ-³²P]ATP (50 µCi/sample; Perkin Elmer, Indianapolis) using T4-polynucleotide kinase (10 U/µl) for 30 min at 37°C. The ³²P-labelled adducts were separated on PEI-cellulose sheets (Machery Nagel, Düren, Germany) by multi-directional thin-layer chromatography (TLC).

The TLC sheets were scanned using Phosphor-Imaging technology (Fujifilm FLA-3000) and DNA adducts levels were calculated from two B[a]PDE-DNA standards with known adducts levels (1 adduct/10⁶ and 1 adduct/10⁷ nucleotides). The major B[a]P-DNA adduct that was used for quantitation purposes in both HepG2 and A549 cells migrated to the same position as the major adduct of the BPDE-DNA adduct standard. In addition, the B[a]PDE-DNA adduct levels were corrected for the amount of DNA in the sample which was assessed by HPLC-UV analysis.

Statistical analysis

Data were expressed as mean ± standard error of the mean (SEM). Statistical analysis was performed using Graphpad Prism 6. To examine differences between the different treatments at each time point, a two-way analysis of variance test (ANOVA) with Bonferroni post hoc multiple comparison was used. Differences were considered to be statistically significant if the p-value was less than 0.05 (P<0.05).

Results

Activity of β -glucuronidase in lung and liver tissues of mice

Intranasal exposure of the mice to LPS resulted in an approximately 2-fold induction of β -glucuronidase activity in lung tissue compared to control (**Table 1**). Similarly, when LPS treatment was combined with B[a]P treatment, a significant 1.5-fold higher activity of β -glucuronidase was observed compared with the B[a]P-treated group without LPS. In addition, bronchoalveolar lavage fluid (BAL fluid) was collected and the β -glucuronidase activity in BAL fluid was lower than in the tissues, but LPS treatment did result in a significant increase of β -glucuronidase when compared to animals that were not treated with LPS (*i.e.* irrespective of B[a]P exposure). On the other hand, in liver tissue a 1.2-fold lower activity of β -glucuronidase was observed in LPS-treated animals when compared to control and B[a]P treated mice, respectively. These changes in β -glucuronidase activity after LPS treatment were in the range of 2 to 30 U/ml β -glucuronidase, and therefore 4 U/ml was used as biologically relevant dose in the subsequent cell culture experiments (see **Table 1**).

Phenotypes after B[a]P exposure with or without β -glucuronidase

Expression of CYP1A1 and CYP1B1

Exposure to B[a]P significantly induced the expression of *CYP1A1* and *CYP1B1* (**Fig 2a** and **Fig 3a**). In A549 cells, the expression of these genes at 6 hours after exposure was increased 56-fold and 5-fold, respectively. Surprisingly, co-incubation with β -glucuronidase inhibited the induction of expression with approximately 50%-90%. Expression of *CYP1A1* and *CYP1B1* in cells that were treated with β -glucuronidase without additional exposure to B[a]P was also reduced when compared to the expression observed in control cells, but this difference did not reach statistical significance. At 24 hours, the expression of *CYP1A1* and *CYP1B1* was induced by B[a]P to 39-fold and 3-fold, respectively. Surprisingly, the induction of both genes was now strongly increased by the presence of β -glucuronidase (191-fold and 6-fold, respectively). At t=48 hours, the induction of *CYP1A1* and *CYP1B1* in cells that were co-exposed to β -glucuronidase remained higher than in cells that were only exposed to B[a]P. The same pattern of changes in gene expression of *CYP1A1* and *CYP1B1* by β -glucuronidase was observed in the absence of B[a]P,

although less pronounced. Moreover, changes in gene expression of *CYP1A1* and *CYP1B1* were essentially similar in HepG2 cells (**Fig 2b** and **Fig 3b**), but the fold changes that were reached were lower than in A549 cells.

Expression of UGT1A6

As shown in **Fig 4b**, in HepG2 cells the expression of UGT1A6 was 3-fold and 11-fold induced by B[a]P when compared to unexposed cells at 6 and 24 hours after exposure, respectively. Addition of β -glucuronidase inhibited UGT1A6 expression after 6 hours when compared to cells treated with B[a]P alone. However, a significant increase in UGT1A6 expression was observed 24 hours after exposure (61-fold and 6-fold compared to unexposed cells and only B[a]P-treated cells, respectively). At 48 hours, the expression of UGT1A6 remained enhanced by B[a]P and B[a]P with additional β -glucuronidase, but the induction levels were three times lower (9-fold and 28-fold, respectively). In addition, a similar pattern of changes in gene expression of UGT1A6 were shown by β -glucuronidase in the absence of B[a]P.

On the other hand, in A549 cells, there was no significant induction or inhibition of UGT1A6 expression by B[a]P nor by β -glucuronidase, but at t=48 hours, all treatments induced the expression of UGT1A6 compared to unexposed cells (see **Fig 4a**).

B[a]P and its metabolites

Extracellular B[a]P metabolite (B[a]P-7,8-diol and B[a]P-9,10-diol) concentrations of A549 cells that were exposed to 1 μ M B[a]P with β -glucuronidase (10 μ g/ml) for 6 hours significantly decreased, when compared to A549 cells that were exposed to B[a]P only (**Fig 5**, right column). However, when cells were exposed for 24 hours, the concentrations of extracellular B[a]P metabolites in the medium of β -glucuronidase-treated cells significantly increased and were higher than in A549 cells that were exposed to B[a]P alone. At 48 hours, the extracellular B[a]P metabolite levels were below the detection limit in both treatments (presence or absence of β -glucuronidase).

The results were essentially similar for HepG2 cells (**Fig 5**, left column). However, the initial difference at t=6 hours was more pronounced, and statistically significant for B[a]P-7,8-diol and B[a]P-9,10-diol.

Moreover, the concentration of unmetabolized B[a]P in the medium showed similar time-dependent patterns in both cell lines (**Fig 5 g and h**). The concentration of unmetabolized B[a]P gradually declined with time. However, after 24 hours of exposure, the concentration of the parent compound in A549 cells was about 3-fold higher ($P<0.05$) in the samples that included β -glucuronidase than in cells without β -glucuronidase. For HepG2 cells, the concentration of B[a]P in the presence of β -glucuronidase was approximately 2-fold higher than in samples that were treated with B[a]P alone. For both cell lines, B[a]P was almost fully metabolized 48 hours after exposure.

B[a]P-DNA adducts level

B[a]P exposure resulted in a time-dependent increase of B[a]P-DNA adduct levels in both cell lines (**Fig 6**). However, the presence of β -glucuronidase altered the kinetics in which DNA adducts were formed: in A549 cells, at $t=6$ and $t=24$ hours, DNA adduct levels were initially lower in cells that were treated with B[a]P and β -glucuronidase. However, a strong increase in DNA adduct levels from 5 adducts per 10^7 nucleotides at $t=24$ hours to 65 adducts per 10^7 nucleotides at 48 hours ($P<0.0001$), was found in β -glucuronidase-treated A549. This was not the case in the samples with B[a]P only. Consequently, at $t=48$ hours, B[a]P-DNA adduct levels were 1.4-fold higher in cells that were treated with β -glucuronidase compared to treatment with B[a]P only. In HepG2 cells, the presence of β -glucuronidase resulted in 1.5-fold, 2-fold and 1.6-fold higher levels of B[a]P-DNA adducts at 6, 24 and 48 hours, respectively, compared to the samples that were treated with B[a]P only ($P<0.05$ at 48 hours).

Potential mechanisms

Can β -glucuronidase interact with B[a]P to prevent B[a]P from entering the cells?

Since B[a]P metabolism seems to be delayed, we studied whether B[a]P could temporarily bind to β -glucuronidase, which could prevent B[a]P from entering the cell. We assumed that binding of B[a]P to β -glucuronidase would interfere with β -glucuronidase activity. Therefore, we assessed the β -glucuronidase hydrolysis of 4MUGIA in the presence B[a]P (**Fig 7**). 10 $\mu\text{g/ml}$ of β -glucuronidase and 1 μM B[a]P were mixed with different concentration of substrate (4MUGIA) at 37°C for 10 hours. The addition of B[a]P lowered the changes in fluorescence units per hours

compared to control. With increasing concentrations of 4MUgIA, the difference of the Δ fluorescence/hour between these two groups became larger. This difference was significant at 100 μ M ($p<0.01$), 250 μ M ($p<0.05$) and 500 μ M ($p<0.001$) of 4MUgIA. The largest difference was observed at the highest concentration of 4MUgIA and was approximately 18% lower than control. In addition, we performed Michaelis-Menten equation to determine the V_{max} and K_m for both reactions. Although the K_m in both reactions is same (0.07 ± 0.01 μ M and 0.07 ± 0.01 μ M for control and B[a]P treated, respectively), the V_{max} in control is significantly larger than in the B[a]P treated sample ($p<0.05$). Therefore, we concluded that there is a noncompetitive inhibition reaction.

Is the inhibitory effect of β -glucuronidase on CYP1A1 expression dependent on its activity?

In order to gain further insight into the role of β -glucuronidase activity in influencing *CYP1A1* expression, an β -glucuronidase inhibitor (D-saccharic acid 1,4-lactone monohydrate) was added to the B[a]P and β -glucuronidase incubations in A549 cells (**Fig 8**). A concentration of 100 μ M fully inhibited β -glucuronidase activity (**Fig 8a**). As shown in **Fig 8b**, the presence of this inhibitor in incubations with B[a]P and β -glucuronidase did not change the inhibitory effect of β -glucuronidase on *CYP1A1* expression (0.13 ± 0.03 and 0.08 ± 0.01 in the absence or presence of inhibitor, respectively) at $t=6$ h. On the contrary, at $t=24$ h, the presence of this inhibitor significantly lowered *CYP1A1* expression when compared to cells that were exposed to B[a]P and β -glucuronidase without inhibitor ($p<0.01$).

Involvement of the Insulin-like growth factor 2/ mannose-6-phosphate pathway

Extracellular enzymes like β -glucuronidase are known to bind to the Mannose-6-phosphate (M6P) receptor (Gonzalez-Noriega et al. 2001), which is also known as IGF2 receptor. The IGF2 receptor can be inhibited by high concentrations of M6P (El-Shewy and Luttrell 2009). β -Glucuronidase inhibited the induction of *CYP1A1* expression by B[a]P at 6 hours of incubation to approximately 10% (**Fig 9**), whereas in the presence of M6P, β -glucuronidase was unable to inhibit *CYP1A1* expression at 6 hours. As shown previously, β -glucuronidase significantly induced

CYP1A1 expression at 24 hours when coincubated with B[a]P ($p<0.01$). *CYP1A1* expression was still enhanced with additional M6P at 24 hours, but the changes in expression were less pronounced.

Discussion

It remains to be established how the cellular response to B[a]P is affected by the presence of extracellular β -glucuronidase, which is released during inflammation. In this study, we demonstrated that β -glucuronidase initially (6 hours after exposure) inhibited gene expression of enzymes that are pivotal in B[a]P metabolism, including *CYP1A1*, *CYP1B1* and *UGT1A6*. As a result, subsequent formation of B[a]P metabolites and the formation of B[a]P-DNA adducts was delayed in the presence of β -glucuronidase. However, at 24 hours of exposure, *CYP* expression was significantly enhanced in β -glucuronidase-treated cells, probably because more B[a]P remained unmetabolized and β -glucuronidase converted B[a]P derived glucuronide metabolites into active B[a]P metabolites. The higher concentration of active B[a]P metabolites continued to trigger the Ah-receptor for gene expression of *CYP1A1* (in both cell lines) and *CYP1B1* (in A549 only). Consequently, the formation of active B[a]P metabolites and DNA adducts at $t=24$ hours was further increased in β -glucuronidase-treated cells. Because of this delayed metabolism of B[a]P, DNA adduct levels could accumulate in cells that were treated with β -glucuronidase and peaked at 48 hours after the initial exposure.

A recent study observed that increased B[a]P-DNA adduct levels in mice that were exposed to B[a]P and intranasally instilled with LPS (Arlt *et al.*, 2015). As LPS can induce an inflammatory response and stimulate the release of β -glucuronidase from neutrophils (Basinska and Florianczyk 2003; Ngkelo *et al.* 2012), we determined the β -glucuronidase activity in lung and liver tissues from these mice (Table 1). Indeed, β -glucuronidase activity was significantly enhanced in lung and BAL-fluid of all LPS-treated animals when compared to the control group. On the other hand, we found that the β -glucuronidase activity was significantly decreased in liver tissue 3 days after being intranasally instilled with LPS. LPS treatment results in the recruitment of neutrophils from the liver to the lung, which could explain the lower β -glucuronidase activity in the liver after LPS treatment (Reutershan *et al.* 2005). The significant increase of β -glucuronidase activity in the lung samples by LPS treatment was approximately 4 U/ml, which we also subsequently used in our *in vitro* experiments.

B[a]P is known to bind to the AhR which stimulates its own metabolism by inducing the expression of *CYP1A1* and *CYP1B1* (Spink et al. 2002). *CYP1A1* and *CYP1B1* play an important role in both B[a]P activation and detoxification (Mosserova et al. 2009), and B[a]P metabolites are further detoxified by glucuronidation. β -glucuronidase will hydrolyse glucuronidated B[a]P metabolites and therefore increase the concentration of active B[a]P metabolites (Shimoi and Nakayama 2005). Higher concentrations of these B[a]P metabolites (e.g. B[a]P-7,8-diol and B[a]P-9,10-diol) enhance the expression of *CYP1A1* and *CYP1B1* (Almahmeed et al. 2004; Spink et al. 2008). Therefore, we expected that addition of β -glucuronidase would increase *CYP1A1* and *CYP1B1* expression. However, we found that β -glucuronidase inhibited *CYP* expression shortly after B[a]P exposure (*i.e.* 6 hours) in both A549 and HepG2 cells. This initial inhibition of gene expression by β -glucuronidase was independent of β -glucuronidase activity, because the β -glucuronidase inhibitor D-saccharic acid 1,4-lactone monohydrate did not change the results. Several studies report the binding of β -glucuronidase to insulin like growth factor 2 receptors (IGF2R), which are located in the cell membrane (Gonzalez-Noriega and Michalak 2001; Urayama et al. 2004; Vogler et al. 2005). Moreover, IGF2 induced AhR in MCF-7 cells (Tomblin and Salisbury 2014). Therefore, we studied the involvement of IGF2R signaling by adding its inhibitor mannose-6-phosphate (M6P) and we found that the inhibitory effect of β -glucuronidase on *CYP*-expression was also blocked. Therefore, we suggest a connection between IGF2R as receptor of extracellular β -glucuronidase and intracellular AhR-signaling after B[a]P exposure.

In addition, *UGT1A6* can be induced by AhR ligands in order to detoxify reactive B[a]P derivatives (Jin et al. 1993). *UGT1A6* is an important enzyme for the detoxification of B[a]P metabolites and is predominantly located in human liver nuclear membranes (Radomska-Pandya et al. 2002; Zheng et al. 2002). This could explain the different kinetics of expression of *UGT1A6* between the two cell lines in this study. Hence, our mRNA expression data of *UGT1A6* exhibited a similar pattern as *CYP1A1* in HepG2 cells, but not in A549 cells. It is known that lung cells have a lower expression of UGT's than liver cells (Ohno and Nakajin 2009), and therefore it was expected that the effects of β -glucuronidase on the metabolism of B[a]P would be more pronounced in the liver-derived HepG2 cells than in the lung-derived A549 cells. Indeed, B[a]P-DNA adducts were already higher in β -glucuronidase-treated HepG2 cells at t=24 hours, whereas in A549 cells an additional 24 hours was needed for further accumulation of DNA adducts. At the 24 h timepoint, the effects of β -glucuronidase on *CYP* expression was at least partly dependent on β -glucuronidase activity. Thus, we suggest that de-glucuronidated metabolites of B[a]P in combination with the higher concentrations of unmetabolized B[a]P, continued to trigger the Ah-receptor and the subsequent expression of

CYP1A1 and *CYP1B1*. Therefore, our results indicate that there are two possible underlying mechanisms resulting in altered B[a]P metabolism and subsequent DNA adduct formation; 1. changes in gene-expression by IGF2 signalling and 2. de-glucuronidation of glucuronidated metabolites.

Moreover, we studied whether B[a]P could temporarily bind to β -glucuronidase which would postpone B[a]P from entering the cell. Indeed, the presence of B[a]P decreased the capacity of β -glucuronidase to deconjugate 4MUGIA, suggesting an interaction between β -glucuronidase and B[a]P. However, this effect is unlikely to explain the present data, because the changes in β -glucuronidase activity by B[a]P are relatively small (<20%) and in our experiments B[a]P is added in excess. However, this interaction may become more relevant at sites of inflammation and low B[a]P exposure.

Since CYP1A1 is considered to be the major enzyme for the activation and detoxification of B[a]P (Arlt et al. 2008), our measured B[a]P metabolites, including B[a]P-7,8-diol paralleled the pattern of *CYP1A1* expression in both cell lines (HepG2 and A549). In addition, it is known that inhibition of *CYP1A1* decreased B[a]P-DNA adduct formation *in vitro* (Endo et al. 2008), but not *in vivo* (Ma and Lu 2007). However, we showed that with additional β -glucuronidase, the metabolism of B[a]P is delayed, which prolonged the effective exposure of cells to unmetabolized B[a]P, ultimately producing more toxic metabolites instead of excretable derivatives of B[a]P. A study using *Cyp1a1*($-/-$) mice revealed that slower metabolic clearance of B[a]P may indeed lead to greater formation of B[a]P-mediated DNA adducts (Uno et al. 2001).

In summary, in this study we showed that β -glucuronidase alters the cellular response towards B[a]P by changing gene expression of *CYP1A1* in both lung and liver derived cells, ultimately causing higher DNA adduct levels. Moreover, we identified that β -glucuronidase may bind to IGF2R, thereby delaying B[a]P metabolism. This study exemplifies the complexity of the effect of inflammation on B[a]P-induced carcinogenesis, which deserves further attention.

Acknowledgements:

This work was supported by a personal grant of the Chinese Scholarship Council (grant numbers CSC201307720049). Work at King's College London is supported by Cancer Research UK (grant C313/A14329)

and in part by the National Institute for Health Research Health Protection Research Unit (NIHR HPRU) in Health Impact of Environmental Hazards at King's College London in partnership with Public Health England (PHE). The views expressed are those of the authors and not necessarily those of the National Health Service, the NIHR, the Department of Health or PHE.

Conflict of interest

The authors declare that there are no conflicts of interest in this study.

References

- Almahmeed T et al. (2004) Benzo[a]pyrene phenols are more potent inducers of CYP1A1, CYP1B1 and COX-2 than benzo[a]pyrene glucuronides in cell lines derived from the human aerodigestive tract *Carcinogenesis* 25:793-799 doi:10.1093/carcin/bgh078
- Arlt VM et al. (2015) Pulmonary Inflammation Impacts on CYP1A1-Mediated Respiratory Tract DNA Damage Induced by the Carcinogenic Air Pollutant Benzo[a]pyrene *Toxicological sciences : an official journal of the Society of Toxicology* doi:10.1093/toxsci/kfv086
- Arlt VM et al. (2008) Metabolic activation of benzo[a]pyrene in vitro by hepatic cytochrome P450 contrasts with detoxification in vivo: experiments with hepatic cytochrome P450 reductase null mice *Carcinogenesis* 29:656-665 doi:10.1093/carcin/bgn002
- Bartholome R et al. (2010) Deconjugation kinetics of glucuronidated phase II flavonoid metabolites by beta-glucuronidase from neutrophils *Drug Metab Pharmacokinet* 25:379-387
- Basinska A, Florianczyk B (2003) Beta-glucuronidase in physiology and disease *Ann Univ Mariae Curie Skłodowska Med* 58:386-389
- Borm PJ, Knaapen AM, Schins RP, Godschalk RW, Schooten FJ (1997) Neutrophils amplify the formation of DNA adducts by benzo[a]pyrene in lung target cells *Environmental health perspectives* 105 Suppl 5:1089-1093
- Colotta F, Allavena P, Sica A, Garlanda C, Mantovani A (2009) Cancer-related inflammation, the seventh hallmark of cancer: links to genetic instability *Carcinogenesis* 30:1073-1081 doi:10.1093/carcin/bgp127
- El-Shewy HM, Luttrell LM (2009) Insulin-like growth factor-2/mannose-6 phosphate receptors *Vitam Horm* 80:667-697 doi:10.1016/S0083-6729(08)00624-9
- Endo K et al. (2008) Inhibition of aryl hydrocarbon receptor transactivation and DNA adduct formation by CYP1 isoform-selective metabolic deactivation of benzo[a]pyrene *Toxicol Appl Pharmacol* 230:135-143 doi:10.1016/j.taap.2008.02.009
- Fridlender ZG, Albelda SM (2012) Tumor-associated neutrophils: friend or foe? *Carcinogenesis*
- Gonzalez-Noriega A, Michalak C (2001) Mannose 6-phosphate-independent endocytosis of beta-glucuronidase. II. Purification of a cation-dependent receptor from bovine liver *Biochim Biophys Acta* 1538:152-161
- Gonzalez-Noriega A, Michalak C, Cruz-Perez JR, Masso F (2001) Mannose 6-phosphate-independent endocytosis of beta-glucuronidase by human fibroblasts. I. Evidence for the existence of a membrane-binding activity *Biochim Biophys Acta* 1538:141-151
- Gungor N, Haegens A, Knaapen AM, Godschalk RW, Chiu RK, Wouters EF, van Schooten FJ (2010a) Lung inflammation is associated with reduced pulmonary nucleotide excision repair in vivo *Mutagenesis* 25:77-82
- Gungor N et al. (2010b) Genotoxic effects of neutrophils and hypochlorous acid *Mutagenesis* 25:149-154 doi:gep053 [pii]
- 10.1093/mutage/gep053
- Gungor N, Pennings JL, Knaapen AM, Chiu RK, Peluso M, Godschalk RW, Van Schooten FJ (2010c) Transcriptional profiling of the acute pulmonary inflammatory response induced by LPS: role of neutrophils *Respir Res* 11:24 doi:1465-9921-11-24 [pii]
- 10.1186/1465-9921-11-24
- Hoffmann D, Hoffmann I (1997) The changing cigarette, 1950-1995 *J Toxicol Environ Health* 50:307-364 doi:10.1080/009841097160393
- Jin CJ, Miners JO, Burchell B, Mackenzie PI (1993) The glucuronidation of hydroxylated metabolites of benzo[a]pyrene and 2-acetylaminofluorene by cDNA-expressed human UDP-glucuronosyltransferases *Carcinogenesis* 14:2637-2639
- Kim JH, Stansbury KH, Walker NJ, Trush MA, Strickland PT, Sutter TR (1998) Metabolism of benzo[a]pyrene and benzo[a]pyrene-7,8-diol by human cytochrome P450 1B1 *Carcinogenesis* 19:1847-1853
- Knaapen AM, Gungor N, Schins RP, Borm PJ, Van Schooten FJ (2006) Neutrophils and respiratory tract DNA damage and mutagenesis: a review *Mutagenesis* 21:225-236 doi:gel032 [pii]
- 10.1093/mutage/gel032
- Krais AM et al. (2015) The impact of p53 on DNA damage and metabolic activation of the environmental carcinogen benzo[a]pyrene: effects in Trp53(+/+), Trp53(+/-) and Trp53(-/-) mice *Arch Toxicol* doi:10.1007/s00204-015-1531-8

- Kucab JE, van Steeg H, Luijten M, Schmeiser HH, White PA, Phillips DH, Arlt VM (2015) TP53 mutations induced by BPDE in Xpa-WT and Xpa-Null human TP53 knock-in (Hupki) mouse embryo fibroblasts *Mutat Res* 773:48-62 doi:10.1016/j.mrfmmm.2015.01.013
- Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method *Methods* 25:402-408 doi:10.1006/meth.2001.1262
- Ma Q, Lu AY (2007) CYP1A induction and human risk assessment: an evolving tale of in vitro and in vivo studies *Drug Metab Dispos* 35:1009-1016 doi:10.1124/dmd.107.015826
- Moseroova M, Kotrbova V, Aimova D, Sulc M, Frei E, Stiborova M (2009) Analysis of benzo[a]pyrene metabolites formed by rat hepatic microsomes using high pressure liquid chromatography: optimization of the method *Interdiscip Toxicol* 2:239-244 doi:10.2478/v10102-009-0024-0
- Ngkelo A, Meja K, Yeadon M, Adcock I, Kirkham PA (2012) LPS induced inflammatory responses in human peripheral blood mononuclear cells is mediated through NOX4 and Gialpha dependent PI-3kinase signalling *J Inflamm (Lond)* 9:1 doi:10.1186/1476-9255-9-1
- Ohno S, Nakajin S (2009) Determination of mRNA expression of human UDP-glucuronosyltransferases and application for localization in various human tissues by real-time reverse transcriptase-polymerase chain reaction *Drug Metab Dispos* 37:32-40 doi:10.1124/dmd.108.023598
- Radomska-Pandya A, Pokrovskaya ID, Xu J, Little JM, Jude AR, Kurten RC, Czernik PJ (2002) Nuclear UDP-glucuronosyltransferases: identification of UGT2B7 and UGT1A6 in human liver nuclear membranes *Arch Biochem Biophys* 399:37-48 doi:10.1006/abbi.2001.2743
- Reutershan J, Basit A, Galkina EV, Ley K (2005) Sequential recruitment of neutrophils into lung and bronchoalveolar lavage fluid in LPS-induced acute lung injury *Am J Physiol Lung Cell Mol Physiol* 289:L807-815 doi:10.1152/ajplung.00477.2004
- Schults MA, Oligschlaeger Y, Godschalk RW, Van Schooten FJ, Chiu RK (2013) Loss of VHL in RCC Reduces Repair and Alters Cellular Response to Benzo[a]pyrene *Front Oncol* 3:270 doi:10.3389/fonc.2013.00270
- Schults MA, Sanen K, Godschalk RW, Theys J, van Schooten FJ, Chiu RK (2014) Hypoxia diminishes the detoxification of the environmental mutagen benzo[a]pyrene *Mutagenesis* 29:481-487 doi:10.1093/mutage/geu034
- Shacter E, Weitzman SA (2002) Chronic inflammation and cancer *Oncology (Williston Park)* 16:217-226, 229; discussion 230-212
- Shimada T, Guengerich FP (2006) Inhibition of human cytochrome P450 1A1-, 1A2-, and 1B1-mediated activation of procarcinogens to genotoxic metabolites by polycyclic aromatic hydrocarbons *Chem Res Toxicol* 19:288-294 doi:10.1021/tx050291v
- Shimoi K, Nakayama T (2005) Glucuronidase deconjugation in inflammation *Methods Enzymol* 400:263-272 doi:10.1016/S0076-6879(05)00015-7
- Shimoi K, Saka N, Nozawa R, Sato M, Amano I, Nakayama T, Kinae N (2001) Deglucuronidation of a flavonoid, luteolin monoglucuronide, during inflammation *Drug Metab Dispos* 29:1521-1524
- Spink DC et al. (2002) Induction of CYP1A1 and CYP1B1 in T-47D human breast cancer cells by benzo[a]pyrene is diminished by arsenite *Drug Metab Dispos* 30:262-269
- Spink DC, Wu SJ, Spink BC, Hussain MM, Vakharia DD, Pentecost BT, Kaminsky LS (2008) Induction of CYP1A1 and CYP1B1 by benzo(k)fluoranthene and benzo(a)pyrene in T-47D human breast cancer cells: roles of PAH interactions and PAH metabolites *Toxicol Appl Pharmacol* 226:213-224 doi:10.1016/j.taap.2007.08.024
- Stiborova M et al. (2014) Cytochrome b5 and epoxide hydrolase contribute to benzo[a]pyrene-DNA adduct formation catalyzed by cytochrome P450 1A1 under low NADPH:P450 oxidoreductase conditions *Toxicology* 318:1-12 doi:10.1016/j.tox.2014.02.002
- Tomblin JK, Salisbury TB (2014) Insulin like growth factor 2 regulation of aryl hydrocarbon receptor in MCF-7 breast cancer cells *Biochem Biophys Res Commun* 443:1092-1096 doi:10.1016/j.bbrc.2013.12.112
- Trushin N et al. (2012) Comparative metabolism of benzo[a]pyrene by human keratinocytes infected with high-risk human papillomavirus types 16 and 18 as episomal or integrated genomes *J Carcinog* 11:1 doi:10.4103/1477-3163.92309
- Uno S, Dalton TP, Shertzer HG, Genter MB, Warshawsky D, Talaska G, Nebert DW (2001) Benzo[a]pyrene-induced toxicity: paradoxical protection in Cyp1a1(-/-) knockout mice having increased hepatic BaP-DNA adduct levels *Biochem Biophys Res Commun* 289:1049-1056
- Uppstad H, Ovrebø S, Haugen A, Møllerup S (2010) Importance of CYP1A1 and CYP1B1 in bioactivation of benzo[a]pyrene in human lung cell lines *Toxicol Lett* 192:221-228 doi:10.1016/j.toxlet.2009.10.025

- Urayama A, Grubb JH, Sly WS, Banks WA (2004) Developmentally regulated mannose 6-phosphate receptor-mediated transport of a lysosomal enzyme across the blood-brain barrier *Proc Natl Acad Sci U S A* 101:12658-12663
- van Berlo D et al. (2010) Neutrophil-derived ROS contribute to oxidative DNA damage induction by quartz particles *Free radical biology & medicine* 49:1685-1693 doi:10.1016/j.freeradbiomed.2010.08.031
- Van Schooten FJ et al. (2004) Myeloperoxidase (MPO) -463G->A reduces MPO activity and DNA adduct levels in bronchoalveolar lavages of smokers *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology* 13:828-833
- Van Schooten FJ et al. (1997) 32P-postlabelling of aromatic DNA adducts in white blood cells and alveolar macrophages of smokers: saturation at high exposures *Mutat Res* 378:65-75
- Vogler C et al. (2005) Overcoming the blood-brain barrier with high-dose enzyme replacement therapy in murine mucopolysaccharidosis VII *Proc Natl Acad Sci U S A* 102:14777-14782
- Wohak LE et al. (2014) Carcinogenic polycyclic aromatic hydrocarbons induce CYP1A1 in human cells via a p53-dependent mechanism *Arch Toxicol* doi:10.1007/s00204-014-1409-1
- Young RP, Hopkins RJ, Christmas T, Black PN, Metcalf P, Gamble GD (2009) COPD prevalence is increased in lung cancer, independent of age, sex and smoking history *Eur Respir J* 34:380-386 doi:10.1183/09031936.00144208
- Zheng Z, Fang JL, Lazarus P (2002) Glucuronidation: an important mechanism for detoxification of benzo[a]pyrene metabolites in aerodigestive tract tissues *Drug Metab Dispos* 30:397-403

Legends to Figures

Figure 1

Critical steps of B[a]P activation and UDP-glucuronosyltransferases (UGTs) detoxification. B[a]P is metabolized to hydroxylated B[a]P including B[a]P-7,8-diol and these B[a]P metabolites are further detoxified by UGTs. β -glucuronidases are able to hydrolyze glucuronidated B[a]P metabolites and therefore increase the amount of B[a]P-7,8-diol, consequently leading to more BPDE and DNA adducts formation.

Figure 2

RT-qPCR analysis of gene expression *CYP1A1* in both A549 (a) and HepG2 (b) cells after exposure to β -glucuronidase and/or B[a]P. Cells were exposed to 1 μ M B[a]P with or without 4 U/ml β -glucuronidase and harvested after the times indicated. Cells exposed to DMSO and sodium acetate buffer were used as a vehicle control. All values are given as the means \pm SEM (n=4 per data point). (*p<0.05; **p<0.01; ***p<0.001)

Figure 3

RT-qPCR analysis of gene expression *CYP1B* in both A549 (a) and HepG2 cells (b) after exposure to β -glucuronidase and/or B[a]P. Cells were exposed to 1 μ M B[a]P with or without 4 U/ml β -glucuronidase and harvested after the times indicated. Cells exposed to DMSO and sodium acetate buffer were used as a vehicle control. All values are given as the means \pm SEM (n=4 per data point). (*p<0.05; **p<0.01; ***p<0.001)

Figure 4

RT-qPCR analysis of gene expression *UGT1A6* in A549 (a) and HepG2 (b) cells after exposure to β -glucuronidase and/or B[a]P. Cells were exposed to 1 μ M B[a]P with or without 4 U/ml β -glucuronidase and harvested after the times indicated. Cells exposed to DMSO and sodium acetate buffer were used as a vehicle control. All values are given as the means \pm SEM (n=4 per data point). (*p<0.05; **p<0.01; ***p<0.001)

Figure 5



HPLC fluorescence analysis of B[a]P-7,8-diol, B[a]P-9,10-diol, 3-OH-B[a]P and B[a]P in HepG2 cells (left column) and A549 cells (right column) after exposure to β -glucuronidase and/ or B[a]P. Cells were exposed to 1 μ M B[a]P with or without 4 U/ml β -glucuronidase and cell medium was harvested after the time indicated. Cells exposed to DMSO and sodium acetate buffer were used as a vehicle control. All values are given as the means \pm SEM (n=3 per data point).  without β -glucuronidase,  with β -glucuronidase

Figure 6

32 P-postlabelling analysis of DNA adducts level in HepG2 cells and A549 cells after exposure to B[a]P with or without β -glucuronidase. (a) HepG2 cells were exposed to 1 μ M B[a]P with or without 10 μ g/ml β -glucuronidase and harvest after the times indicated. (b) A549 cells were exposed to 1 μ M B[a]P with or without 4 U/ml β -glucuronidase and harvest after the times indicated. Data are expressed as number of B[a]P-DNA adducts per 10^7 nucleotides (n=5 for HepG2 cells and n=4 for A549 cells, mean \pm SEM). (*p<0.05; ***p<0.001). Representative chromatograms obtained by 32 P-postlabeling in HepG2 cells (c) and A549 cells (d). The adduct spot (arrow) that migrated during 2D-TLC to the same position as the major DNA adduct in a BPDE-DNA adduct standard was quantitated in all samples. Before phosphorimaging of the TLC plates the origin located at the bottom left-hand corner was excised.

Figure 7

The 4MUGIA assay was applied to assess the possibility of B[a]P to bind to β -glucuronidase. A total volume of 140 μ l contained 0.1 M sodium-acetate buffer (pH 5.5), different concentration of 4MUGIA (e.g. 500, 250, 100, 50, 10 and 1 μ M) and 4 U/ml β -glucuronidase with 1 μ l of 200 μ M B[a]P or 1 μ l DMSO. The measurement of fluorescence (Relative Fluorescence Unit (RFU)) was performed for 10 hours at 37°C. B[a]P treated samples were compared with control at each concentration, respectively. (*p<0.05; **p<0.01; ***p<0.001)

Figure 8

(a) 4MUGIA assay was applied to assess the inhibition of β -glucuronidase by D-saccharic acid 1,4-lactone monohydrate. A total volume of 140 μ l contained 0.1 M sodium-acetate buffer (pH 5.5), 2 mM 4MUGIA, 4 U/ml β -glucuronidase and different concentration of D-saccharic acid 1,4-lactone monohydrate (10 μ M, 100 μ M and 500

μM). 100 μM of D-saccharic acid 1,4-lactone monohydrate was used in the following incubation. (b) A549 cells were exposed to 1 μM B[a]P with or without β-glucuronidase, D-saccharic acid 1,4-lactone monohydrate for 6 and 24 hours. Cells exposed to 1 μM B[a]P and sodium acetate buffer was used as a control. All values are given as the means ± SEM (n=4 per data point). (*p<0.05; **p<0.01; ***p<0.001)

Figure 9

RT-qPCR analysis of gene expression *CYP1A1* in A549 cells after exposure to β-glucuronidase and B[a]P with the IGF2R inhibitor M6P. Cells were exposed to 1 μM B[a]P with or without 4 U/ml β-glucuronidase and with or without 100 μM M6P. Cells were harvested after the times indicated. Cells exposed to B[a]P and sodium acetate buffer were used as vehicle control. All values are given as the means ± SEM (n=4 per data point). (*p<0.05; **p<0.01; ***p<0.001)

Table 1 Activity of β -glucuronidase in mouse liver, lung tissues and BAL fluids.

| | Control (U/ml) | LPS (U/ml) | B[a]P (U/ml) | B[a]P&LPS (U/ml) |
|---------------|-------------------|-----------------------------|-----------------|------------------------------|
| Liver cytosol | 27.1 \pm 1.5 | 23.3 \pm 0.4 [*] | 27.6 \pm 0.9 | 22.5 \pm 1.0 ^{**} |
| Lung cytosol | 7.6 \pm 0.8 | 15.4 \pm 1.3 [*] | 7.4 \pm 0.8 | 11.1 \pm 1.3 ^{**} |
| BAL fluids | 2 \pm 0.2 | 2.2 \pm 0.1 [#] | 2.0 \pm 0.1 | 2.2 \pm 0.1 [#] |

^{*} significantly different from control animals (p<0.05)

^{**} significantly different from B[a]P-treated animals (p<0.05)

[#] p<0.05 if both LPS-treated groups were combined when compared to non-LPS-treated animals.





Figure 1

(a)



(b)





Figure 2

(a)



(b)





Figure 3

(a)



(b)




Figure 4




Figure 5




Figure 6





Figure 7

(a)



(b)




Figure 8




Figure 9